

Review

Intracranial Electrophysiology of the Human Default Network

Kieran C.R. Fox,^{1,2,*} Brett L. Foster,³ Aaron Kucyi,^{1,2} Amy L. Daitch,^{1,2} and Josef Parvizi^{1,2,4,*}

The human default network (DN) plays a critical role in internally directed cognition, behavior, and neuropsychiatric disease. Despite much progress with functional neuroimaging, persistent questions still linger concerning the electrophysiological underpinnings, fast temporal dynamics, and causal importance of the DN. Here, we review how direct intracranial recording and stimulation of the DN provides a unique combination of high spatiotemporal resolution and causal information that speaks directly to many of these outstanding questions. Our synthesis highlights the electrophysiological basis of activation, suppression, and connectivity of the DN, each key areas of debate in the literature. Integrating these unique electrophysiological data with extant neuroimaging findings will help lay the foundation for a mechanistic account of DN function in human behavior and cognition.

Toward a Deeper Understanding of the Default Network

Serendipity has played an eminent role in the history of scientific discovery: chance observations and insights are alleged to lie at the root of many major and minor breakthroughs [1]. Every innovation must subsequently be refined and scrutinized, however; every paradigm shift in scientific thinking must be followed by years or decades of rigorous and painstaking ‘normal science’ [2]. The fortuitous discovery of the default mode network [or **default network (DN)**, see [Glossary](#)] some 20 years ago provides a prototypical example [3–5]: the unexpected observation that a consistent set of brain regions showed reduced cerebral blood flow (‘deactivated’) during a variety of cognitive tasks (compared to rest [4,5]) seeded a paradigm shift in our view of the functional architecture of the brain [6–8].

The human DN is now well known as a collection of associative brain regions distributed over the temporal, parietal, and frontal lobes ([Figure 1](#)), readily identifiable by correlated spontaneous fluctuations in the fMRI blood oxygen level-dependent (BOLD) signal. The DN is just one of numerous distributed **intrinsic brain networks** [9–12] identified during the resting state and persistently correlated during a variety of cognitive tasks [13,14] and states of consciousness [15–17]. Despite much progress with neuroimaging [18], persistent questions still linger concerning the electrophysiological underpinnings, fast temporal dynamics, and causal importance of the human DN. We require a fuller understanding of how this canonical brain network contributes to human cognition and behavior, and how its pathological modulation relates to neuropsychiatric disorders. Such an understanding calls for a detailed neuromechanistic account that explains the DN’s internal dynamics and interactions with other intrinsic networks with high anatomical precision and temporal resolution. Moreover, the relationships between DN activity and human subjective experience and behavior need to be explored with causal methods that go beyond mere correlative observations.

Highlights

Understanding of the default network (DN) is rapidly progressing from a relatively coarse picture toward more detailed models, in which network nodes contain functionally heterogeneous neuronal populations with distinctive intra- and internetwork connectivity.

Intracranial electroencephalography (IEEG) is providing unique data about the precise temporal profiles of neuronal populations throughout the DN, and their functional role(s) in the network’s deactivations, activations, intrinsic activity, and interactions with other brain networks.

IEEG is also providing critical corroborative data elucidating the electrophysiological foundations of signals observed with neuroimaging.

IEEG methods are being used not only to passively record but also to causally probe and perturb the DN’s functioning and interactions with other networks.

Intracranial methods are being developed as electroceutical interventions for neuropsychiatric disorders implicating the DN.

¹Department of Neurology and Neurological Sciences, Stanford University, Stanford, CA, USA

²Stanford Human Intracranial Cognitive Electrophysiology Program (SHICEP), Stanford, CA, USA

³Departments of Neurosurgery and Neuroscience, Baylor College of Medicine, Houston, TX, USA

⁴Stanford University School of Medicine, Stanford University, Stanford, CA, USA

*Correspondence: kcrfox@stanford.edu (Kieran C.R. Fox) and jparvizi@stanford.edu (J. Parvizi).




Figure 1. Intracranial Electrophysiology of the Human Default Network. (A) The human default network (DN) as assessed using intrinsic connectivity fMRI during the resting state in 1000 participants [9]. The network is distributed throughout the brain: in the frontal lobe, medial prefrontal cortex (PFC), dorsomedial PFC, and ventrolateral PFC; in the temporal lobe, temporopolar cortex, lateral temporal cortex, and aspects of the medial temporal lobe (MTL); and in the parietal lobe, the posterior cingulate cortex and retrosplenial cortex medially, and angular gyrus laterally. Although there is general consensus on this collection of regions, specific network borders vary depending on methods and modality used, and the notion of what constitutes the DN continues to evolve [55]. Because much of the MTL falls outside DN boundaries using standard methods, as a rule we do not discuss iEEG investigations of the MTL throughout this review unless they bear directly on questions relating to the function of other DN regions (for a review covering many iEEG studies involving the MTL, see [103]). (B) Combining iEEG data from cohorts of patients can provide comprehensive coverage of the brain, including all cortical DN regions (here we show 1955 electrode sites from 16 patients). There are two basic types of iEEG recording [19]: (i) electrocorticography (ECoG), using subdural grids or strips of circular (plate-shaped) electrodes placed on the brain's surface, can record from lateral or medial cortical DN areas. Grids often contain dozens of contacts, allowing coverage of, and simultaneous recording from, a substantial proportion of cortex. (ii) Stereotactic EEG (SEEG) uses cylindrical depth electrodes which penetrate through the skull and brain tissue and can reach deep medial cortical and subcortical structures, including midline DN regions. Each depth electrode typically contains 6 to 10 contacts. For more details, see [19] and Box 1.

Intracranial electroencephalography (iEEG) (Box 1) has much to contribute to this endeavor, by providing direct recording and stimulation data from the human brain. Typically, iEEG is undertaken in patients undergoing monitoring for epilepsy surgery, and involves the placement of electrode grids directly on the cortical surface (see Figure 1 in Box 1), and/or the insertion of depth electrodes which can contact deep cortical and subcortical structures. With high anatomical precision (at the level of **neuronal populations**), high temporal resolution (at the millisecond scale), and high signal-to-noise ratio (at the level of several-fold signal increase) [19], iEEG provides the necessary fine-scale view of the human brain to both advance and refine our knowledge of the DN. Although the tools of iEEG and electrical brain stimulation have been available for decades, recent advances in computational methods have given rise to entirely novel means of charting and perturbing the engagement of local neuronal populations and larger functional networks. For example, gauging the power of the electrophysiological signal in

Glossary

Default network (DN): a collection of associative brain regions that consistently deactivate during externally directed tasks (e.g., visual search), while conversely showing activation during internally directed tasks (e.g., memory retrieval). Like other intrinsic networks, DN regions display robustly correlated hemodynamic activity during resting states. Anatomically, key nodes of the DN include the medial prefrontal cortex (retrosplenial cortex, posterior cingulate cortex), aspects of the medial temporal lobe, inferior lateral parietal cortex (angular gyrus), middle temporal gyrus, and parts of ventrolateral and dorsomedial prefrontal cortex (Figure 1).

Electroceutical interventions: clinical interventions utilizing intracranial electrical stimulation and modulation to ameliorate neurodegenerative (e.g., Parkinson's disease) and psychiatric (e.g., major depression) illness; focal neuromodulation nonetheless has indirect effects on the metabolism and functioning of widespread brain networks.

Envelope: an alternative term for the time varying 'analytic amplitude' of a frequency band limited signal, typically squared to obtain 'power'. A common metric used for correlating task or resting-state activity between intracranial electroencephalography (iEEG) recording sites.

High-frequency broadband (HFB): nonoscillatory, high-frequency (>50 Hz) activity in the cortical field potential; a primary signal of interest in iEEG studies, known to be correlated with the fMRI blood oxygen level-dependent (BOLD) signal as well as averaged neuronal population spiking activity.

High-frequency electrical brain stimulation (hf-EBS): injection of electrical current into brain tissue via intracranial electrodes, typically using square-wave pulses at ≥ 50 Hz and 1–10 mA. Routinely used in clinical mapping sessions to determine localization of function, hf-EBS often elicits subjective experiences or perturbs ongoing cognition and perception, providing a potent tool for causally exploring regional functionality.

Trends in Cognitive Sciences

Box 1. An Overview of the Intracranial Electroencephalography Method

Functional and morphometric MRI have undoubtedly played the starring role in the discovery and subsequent investigation of the default network (DN) [4,8,42,105]. Yet even as the DN continues to occupy the spotlight in much of contemporary cognitive and clinical neuroscience research, noninvasive and correlational neuroimaging methods continue to confront important technological barriers. Limited temporal resolution, poor signal-to-noise ratio, and lack of causal data from neuroimaging have left in their wake many persistent and critical questions that intracranial electroencephalography (iEEG) can make a unique contribution to answering.

The spatial precision of the iEEG signal lies between that of classic local field potentials and scalp EEG [106]. Similar to scalp EEG, iEEG records the brain's endogenously produced electrical potentials, but by bypassing the filtering properties of the meninges, skull, and scalp, iEEG gains a huge increase in signal-to-noise ratio and a widening of the detectable spectrum of neural activity. Moreover, the recorded signal has great anatomical precision, since the inverse problem (i.e., inferring the precise source of recorded signals) is greatly mitigated. The fundamental recorded activity is the cortical field potential, a rich signal thought to represent a summation primarily of synaptic currents and neuron spiking near the electrode [19,106,107]. Although the recorded signal does not reveal the precise spiking of specific neurons (the 'trees'), it does reliably measure the engagement of the 'forest' (population) of neurons (an estimated 200 000–500 000 neurons [19]). iEEG therefore provides mesoscale-level data, uniquely complementary to the cellular level of mapping with single-neuron recordings in animals and the regional/whole-brain level of mapping in humans, with neuroimaging.

iEEG's powerful combination of high anatomical and temporal resolution can contribute unique and hitherto unknown information to the field of human neuroscience. With electrode contacts placed directly on the surface of the cortex (in grids; Figure 1) or deep within brain structures (with cylindrical depth electrodes), unlike single-unit recordings, iEEG can often capture simultaneous recordings across a large mantle of the human brain (up to ~200 electrodes in a single patient). By recording simultaneously from several nodes of a network, iEEG signals can reveal information about functional interactions within and across networks during different stages of neural computation even during complex cognitive, affective, and perceptual tasks requiring human participants. Furthermore, the method allows delivery of a volley of electrical discharges directly to a brain region in awake human participants [93,94], providing the ability to causally perturb subjective experience and behavioral performance, as well as modulate network activity (Box 3) and potentially treat neuropsychiatric disease (Box 4).




Figure 1. Intracranial Electroencephalography Refers to the Invasive Measure of Electrical Brain Potentials via Two Common Methods. (A) Electrocorticography (ECoG) utilizes strips or grids of electrode arrays placed subdurally on the cortical surface. (B) Stereo-electroencephalography utilizes penetrating depth electrode arrays based on stereotactic coordinates for targeting deeper brain structures.

Intracranial

electroencephalography (iEEG): a suite of methods for invasively monitoring the brain's electrical activity and directly stimulating brain tissue. Includes grids and strips of electrodes placed subdurally on the cortical surface (also known as electrocorticography, or ECoG), as well as depth electrodes that penetrate brain tissue and reach deep cortical and subcortical structures (also known as stereotactic EEG, or SEEG).

Intrinsic brain network: a set of spatially segregated brain regions displaying interregional correlated spontaneous activity (on the scale of minutes and longer) that is persistent across cognitive and conscious states (e.g., tasks and rest, waking and sleep).

Neuronal population: the group of nearby neurons (~200 000–500 000) thought to make the main contribution to the local signal recorded from a given intracranial electrode contact.

the **high-frequency broadband (HFB)** range (>50 Hz, also known as 'high gamma'; see Box 2) has provided a new metric of cortical activation that correlates well with both fMRI BOLD signals [20,21] and averaged neuronal spiking activity [22], providing a potent means of replicating and extending findings from human neuroimaging and animal neurophysiology, respectively (Box 3).

Box 2. Understanding High-Frequency Activity in the Brain

The high-frequency broadband (HFB) signal is currently interpreted as a reflection of a non-oscillatory broadband signal [146,147]. Importantly, HFB differs from pathological high-frequency oscillations (HFOs) that are seen in epileptic recording sites. Unlike HFB signals, HFOs are associated or coupled with interictal epileptiform discharges and/or pathological background activity at the recording site and can be present across several adjacent electrodes [148,149] (Box 1).

A growing body of evidence suggests that high-frequency activity is a reliable electrophysiological correlate of underlying population spiking activity generated by thousands of neurons [21,150–153]. HFB signals also correlate with hemodynamic signals detected with fMRI [20–22,154,155]. Thus, an increase in the HFB power in a recording site represents the local engagement of the cortical tissue underneath or around the electrode. There is strong evidence that, unlike slow oscillatory activity, the HFB signal has a remarkably localized anatomical precision and originates from the cortical tissue immediately around or underneath the recording electrode; for human intracranial electroencephalography (iEEG) evidence, see [28,30,47,156–159], and for direct measures from non-human primates, see [151–154].

There is evidence from non-human brains that the number of cells contributing to the high-frequency signal may be as little as ~16% of neurons sampled by a given electrode [160] and that the source of the high-frequency signal is often within several hundreds of micrometers from the electrode tip [152,161,162]. Given the specifics of iEEG electrodes and recording parameters, our assumption is that the iEEG signal is rooted in the activity of a diverse population of cells in the millimeter space (for a recent, in-depth review, see [19]).

This Review provides an accessible synthesis of the most recent iEEG findings related to the DN for researchers both in and outside the field of human intracranial electrophysiology. We aim to promote crosstalk among disciplines and explore how iEEG findings can help transform our understanding of the DN. In the text that follows, we synthesize the DN iEEG literature pertinent to: (i) task-induced deactivations, (ii) task-induced activations, (iii) intra- and internetwork connectivity, and (iv) the causal effects of electrical stimulation.

Box 3. Advantages of Combining Intracranial Electroencephalography with Other Modalities

There are many applications of combining intracranial electroencephalography (iEEG) with other methods, particularly fMRI. Here we highlight several examples, but the possibilities of such multimodal investigations are still just beginning to be explored.

Providing Functional Context for Electrode Localization: Although the anatomical location of intracranial electrodes can be reconstructed with relatively high precision, functional brain networks vary considerably across individual neuroanatomy [108,109]. Anatomical landmarks alone are therefore at best a coarse indicator of regional function, but resting-state fMRI scans can provide relatively reliable estimates of subject-specific brain networks [110,111], allowing particular electrodes to be assigned with some confidence to individually specified functional networks [87,111,112].

Putting Novel iEEG-Elicited Effects into Broader Context: While iEEG can discover novel effects, its sparse coverage of the brain often renders it uninformative about the whole-brain relevance of these unique findings. fMRI can supplement the paucity of recording sites in iEEG and provide complementary whole-brain data that clarifies and corroborates iEEG findings (e.g., [112]).

Understanding Effects of Focal Electrical Stimulation in Brain Disease: With the increasing use of intracranial methods to treat neurological and psychiatric disease, combined pre- and postintervention positron emission tomography and fMRI scanning has already been employed to elucidate how focal **electroceutical interventions** affect functioning of the default network [113–115]. Neuroimaging will continue to play a key role in corroborating and understanding the brain-wide and long-term effects of targeted intracranial stimulation (Box 4).

Simultaneous Multimodal Investigations: Progress is being made in developing ways of safely combining iEEG with both fMRI [116,117] and magnetoencephalography (MEG) [118,119] in humans, which could open up many new possibilities, for instance directly observing the whole-brain effects of focal electrical stimulation [120].

Electrophysiological Correlates of Task-Induced Default Network Deactivations


The first finding from neuroimaging suggesting the DN was a coordinated functional system was the observation of reduced cerebral blood flow in DN regions during engagement in externally directed tasks, such as visual search [4,5], compared with a resting or baseline state. Some of the earliest intracranial investigations therefore explored whether parallel electrophysiological markers of deactivation could be observed with iEEG (Figure 2A–C). For iEEG data, ‘deactivation’ would be reflected in reductions in power of high-frequency activity (such as HFB) relative to the pre-stimulus period (typically, 100–200 ms prior to the onset of a given stimulus). Beginning in 2008, researchers began to note suppression of HFB power in response to externally directed tasks in isolated DN regions, such as ventrolateral [23] (20–150 Hz) and rostromedial [24] prefrontal cortex (PFC) (50–150 Hz). Subsequent studies reported similar deactivations in HFB activity while recording from multiple DN regions simultaneously [25–27]. Critically, some studies have explored the various cognitive and behavioral correlates of electrophysiological DN deactivations [27,28], and parallel work has confirmed that anatomical sites of iEEG deactivation are indeed localized within subject-specific DN boundaries as identified with resting-state fMRI [29,30].

Collectively, these investigations have revealed essential information about the DN specifically and the fMRI BOLD signal more generally. First, these data have helped clarify the neurophysiological contributions to fMRI BOLD signal decreases. Early concerns had been raised that apparent deactivations in neuroimaging data were of non-neural origin, for instance, caused by cardiac or respiratory artifacts [31] or vascular ‘stealing’ [32]. Task-induced suppression of iEEG HFB activity, a correlate of averaged neuronal firing, provides compelling evidence that DN deactivations seen in neuroimaging studies have a firm grounding in the electrophysiological activity of neuronal populations within DN regions (Figure 2A), a conclusion further supported by single-neuron data from non-human primates [33].

Second, HFB suppression in various DN areas is observed in response to a wide variety of externally directed tasks (as with neuroimaging [5]), including visual search [27], reading [23,26], a backward-masking visual categorization task [34], the Navon ‘global versus local’ attention task [26], and arithmetic calculation [28–30,35]. These findings support the conclusion from neuroimaging studies that DN deactivations are a common, domain-general response to the demands of externally directed attention.

Third, not all DN areas deactivate at the same time [26,27] (Figure 2B; but for an exception, see [34]). The finding of subtle differences in mean timing of iEEG deactivations across DN nodes highlights an important contribution made possible by the high temporal resolution of iEEG. Indeed, the ability to temporally dissociate responses within the DN is a critical domain of progress uniquely served by iEEG (see ‘Electrophysiological Correlates of Intra- and Internetwork Interactions,’ below).

Finally, the timing and magnitude of HFB suppression have functional significance. The duration of HFB suppression correlates with reaction time [27,28] (consistent with magnitude-reaction time correlations with fMRI [36]). Additionally, both the duration and magnitude of HFB deactivations are associated with task complexity, with more complex tasks inducing longer and larger iEEG deactivations [27] (Figure 2C) (a finding observed early on with fMRI [5,37]). Both findings suggest that deactivation represents not merely the relative quiescence of a network not participating in a given task, but instead might be an active process of functional



Trends in Cognitive Sciences

Figure 2. Unique Contributions to Understanding Default Network Deactivations and Activations from Intracranial Electroencephalography. (A) Default network deactivation in response to a visuomotor task consists largely of rapid suppression of power in higher frequency ranges (>30 Hz). Hot colors indicate increased power, and cool colors decreased power, at given frequencies. These, and similar findings from other studies, constitute a major advance in understanding the electrophysiological basis of default network (DN) deactivations observed with neuroimaging. (B) Timing of deactivation matters: whereas neuroimaging findings might suggest that the entire DN deactivates simultaneously, intracranial electroencephalography (IEEG) shows that there are subtle (yet probably important) timing differences in the onset of higher frequency 'gamma band deactivations' (GBD) in response to the presentation of complex visual task stimuli. (C) The duration of deactivations is also of functional significance: a more difficult visual search task elicited significantly longer suppressions of high-frequency power in default regions. (D) Certain neuronal populations in the posteromedial cortex (PMC) are significantly activated (electrodes with red fill color) by a variety of internally directed cognitive tasks; (E) eliciting power increases specifically at higher frequencies (hot colors indicate increased power, cool colors decreased power). (F) The mean magnitude of PMC activations in the high-gamma (HG) band (70–180 Hz) can differentiate between conditions, with the largest activations for the most episodic/memory-like judgments. By contrast, significant suppression of high-gamma power is evident during an externally directed math task (i.e., judging the accuracy of simple arithmetic equations); ** indicates activity is significantly different from all other conditions ($P < 0.01$, corrected). (G) Intraregional functional heterogeneity within the PMC. Certain populations show preferential activation in response to memory retrieval, others are selectively active during rest periods, and yet others show mixed patterns. Default regions often treated as functionally homogeneous in neuroimaging investigations in fact contain populations with very diverse response profiles. (A) Reproduced, with permission, from [34]; (B,C) reproduced, with permission, from [27]; and (D–F) reproduced, with permission, from [28]; and (G) adapted, with permission, from [52]. Abbreviations: LTC, lateral temporal cortex; MPFC, medial prefrontal cortex; TPJ, temporoparietal junction; VLPFC, ventrolateral prefrontal cortex.

suppression important for successful task performance. Increasing task complexity/difficulty also reliably leads to reductions in self-reported mind-wandering and other internally directed thought processes (e.g., [38,39]); whether or not these psychological effects are directly related to differential duration and magnitude of deactivation in default areas is an important question that should be explored in future research (cf., [37]).

Electrophysiological Correlates of Task-Induced Default Network Activations

Whereas externally directed cognitive tasks often evoke DN deactivation, DN activation (i.e., relative increases of cerebral blood flow or BOLD signal strength compared with the control condition) is linked to many forms of internally directed cognition [40,41] such as self-referential thinking [42], mind-wandering [43,44,163], simulation of the future [45], and remembrance of the past [46]. The unique strengths of iEEG are helping to disentangle the complex functionality and electrophysiology of DN activation by providing detailed information about the time course, magnitude, and selectivity of signals from recording sites throughout the network. Collectively, investigations of DN activation have allowed us to reach several broad conclusions about the temporal dynamics of, and role(s) played by, different regions and neuronal populations.

First, the electrophysiology of DN activations comprises increases in HFB power (compared with pre-stimulus baseline) concomitant with decreases in lower frequency ranges, such as alpha and theta [29,30] (Figure 2E), mirroring results from intracranial investigations of motor and sensory cortices [47]. This finding complements the spectral changes that characterize DN deactivations, which typically show the opposite pattern of power change [27,34] (Figure 2A), and contrasts with intrinsic resting state DN activity, where theta oscillations predominate [48].

Second, similar to the breadth of internally directed processes that activate the DN in neuroimaging [49,50], a variety of internally directed tasks increase HFB power in DN neuronal populations [28–30,51,52] (Figure 2F). Increased high-frequency (30–180 Hz) power was first reported [29] in the posterior cingulate cortex (PCC) and retrosplenial cortex (RSC) during self-referential judgments (consistent with fMRI [53]), with numerous iEEG studies having since replicated these observations in these and other default areas [28,30,51,52].

Third, the magnitude of DN activations has functional significance (as it does for deactivations). Specific neuronal populations (i.e., electrode sites) showing larger HFB power increases for a given type of internally directed cognition (such as episodic memory recall) tend to be more active for similar kinds of mentation, such as self-referential judgments [28]. Activation magnitude can also distinguish between different types of internally directed cognition. For instance, a recent study [28] found a clear tuning of HFB activations for self-referential judgments in PCC and RSC: larger responses were elicited by more episodic/autobiographical judgments (e.g., 'I ate fruit yesterday'), compared with more general semantic (e.g., 'I eat fruit often') or personal judgments (e.g., 'I am kind') (Figure 2F). Moreover, PCC/RSC neuronal populations more active for autobiographical judgments also showed larger deactivations during an externally directed attentional task (arithmetic calculation) [28].

Fourth, not all populations within DN areas respond in the same way (Figure 2G), suggesting that large-scale parallels between DN activation across fMRI and iEEG represent only a coarse picture of averaged signals in a given region. Although DN regions, and even the entire network, are often treated as functionally homogenous in the neuroimaging literature, some pioneering efforts at finer functional-anatomic fractionations have been undertaken in recent years [54–57]. These fMRI studies hint at the possibility that particular DN regions, too, might contain functionally heterogeneous neuronal populations. In keeping with this, a study [29] using direct recordings from the DN hub in the posteromedial cortex (PMC) found only a small subset of recording sites displaying high-frequency (30–180 Hz) power increases to self-referential judgments. A more recent study [52] involved a larger cohort of 13 individuals who alternated between evaluating autobiographical memory and arithmetic statements, and staring at a fixation cross at the center of a dark blank screen ('fixation rest'). Recording from multiple sites within each individual patient's PMC revealed clear functional heterogeneity across three distinct populations of neurons within this DN hub.

Anatomically distinct clusters of neuronal populations in the PMC showed activations (in the HFB range) during either memory processes or fixation rest, and within-subject analysis showed that rest-active sites were (on average) located more dorsally than memory-active sites. Such spatially overlapping but functionally heterogeneous populations (Figure 2G) can only be appreciated with a method that offers a unique combination of subject-specific anatomical precision, high signal-to-noise ratio, and simultaneous recordings with tens of electrodes across a cortical region (Box 1). Moreover, there were subtle differences in the temporal domain as well. Some populations showed fast, time-locked activations at the beginning of fixation rest trials (i.e., 'switch sites'). These neuronal populations might help shift the brain from an externally directed attentional state to a more internally focused one, as participants become free to explore their own thoughts and surroundings (cf. [58,59]). Conversely, this fast, time-locked, and seemingly automatic activity following the switch from externally focused attention to fixation rest is unlikely to reflect the re-emergence of spontaneous thoughts (as the time of onset of HFB activity is too quick for any spontaneous thought to occur). Other clusters of PMC neurons showed a slow, temporally jittered rise of activity during fixation rest. Such variable and delayed temporal dynamics suggest that these populations of PMC neurons could be engaged in the arising of stimulus-independent spontaneous cognition, which is unlikely to occur with consistent timing across trials [60]. Finally, a third type of neuronal population in the PMC showed time-locked activations during the memory condition, and often deactivated during the math condition; the relatively late (>300 ms) recruitment of these populations suggests that the PMC is likely engaged after the medial temporal lobe (MTL).

Although high-resolution/long-duration single-subject recordings and multi-voxel pattern analysis with fMRI have begun to provide considerably greater spatial precision, the distinct temporal profiles described above would be impossible to isolate with neuroimaging methods.

Electrophysiological Correlates of Intra- and Internetwork Interactions

Network neuroscience views the brain as functioning via widely dispersed but interconnected regions coordinating their activity across multiple timescales [61,62]. Fine temporal resolution methods are therefore needed to track how neocortical dynamics are modulated at both fast (>40 Hz) and slow (< 1 Hz) frequencies. iEEG is now beginning to furnish detailed data about fast temporal dynamics to complement slower-timescale data from neuroimaging. These new data can advance our understanding of intra- and internetwork interactions in several ways. First, iEEG can explore the electrophysiological correlates of intrinsic activity and connectivity and how these relate to network models from neuroimaging. Second, iEEG can explore the temporal priority of network recruitment during DN activation by examining the time course of activity throughout the brain. Third, direct electrical stimulation can be utilized to probe the directionality and time course of the DN's intra- and internetwork interactions using cortico-cortical evoked potentials (CCEPs). We discuss each approach sequentially in the following sections.

Intrinsic Activity of the Default Network

Scalp-based EEG has a long history of investigating spontaneous fluctuations in the brain's electrical activity and identifying canonical oscillations reliably associated with various parts of the brain [63,64]. Can similar observations be made for the DN? A study in 2012 [48] was the first to report that the PMC displayed strong intrinsic theta band oscillations (~4–5 Hz peak) at rest, in contrast to nearby visual areas, which instead showed canonical alpha band oscillations (~8–10 Hz peak). Analysis of cross-frequency coupling revealed that HFB (70–180 Hz) amplitude at rest was coupled with the phase of ongoing theta oscillations in the PMC (i.e., phase-amplitude coupling), suggesting that patterns of intrinsic activity in DN areas (at least PMC) are readily distinguishable from other brain regions [48]. This study demonstrated that the PMC shows patterns of intrinsic oscillatory activity that are distinct from nearby sensorimotor brain

regions, yet similar to distal areas like the MTL [65], with which PMC likely cooperates to instantiate memory recall and other forms of internally directed cognition [48]. More speculatively, the authors noted that theta/HFB coupling displayed slow modulation (<1 Hz), providing a potential link between local population coordination and slower intrinsic connectivity dynamics [48]. These findings were an important step toward identifying a potential oscillatory 'signature' for the DN, but did not test if these motifs are correlated across multiple DN regions.

Intrinsic Connectivity of the Default Network

The defining feature of intrinsic brain networks identified with BOLD fMRI is the spontaneous correlation of slow (<0.1 Hz) signal fluctuations [66,67]. Given that iEEG signals can vary on a much faster timescale than fMRI BOLD signals, how can these different timescales be linked? One common solution is to examine the low-frequency components of fast activity by extracting the slow fluctuations (i.e., low-pass filtering) of the amplitude **envelope** of a frequency band of interest (e.g., the HFB range [68,69]).

To date, this approach has proven to be a fruitful assay of fMRI-like connectivity patterns [30]. For instance, taking such an approach to spontaneous iEEG connectivity, it has been shown [68] that slow spontaneous fluctuations (<0.1 Hz) in high-gamma (40–100 Hz) envelope signals were highly correlated between the left and right auditory cortex. Early on, researchers working with non-human primates had proposed that the anatomical selectivity of such correlations might mimic functional connectivity patterns seen in fMRI [69], a hypothesis echoed by others [68]. A number of follow-up studies have confirmed this prediction, showing consistent spatial patterns of BOLD fMRI and iEEG-envelope functional connectivity within subjects, including throughout many nodes of the DN [30,70–72].

Employing a related yet analytically distinctive approach, intracranial recordings from medial PFC and PCC showed that these spatially distributed DN regions exhibited coherent activity peaking at ~0.017 Hz at higher frequencies (65–110 Hz) (i.e., envelope spectral coherence) [73]. This frequency of interregional coherence was almost identical to the peak power observed in intrinsic BOLD signal fluctuations recorded from the same patients [73]. Using methods more akin to earlier work [68], more recent research [30] showed that correlation patterns over medial and lateral parietal cortex using iEEG consistently identified DN subregions, specifically the PCC/RSC and angular gyrus (AG). These patterns of iEEG connectivity, using slow (<1 Hz) fluctuations in HFB amplitude, showed high spatial correlation with fMRI connectivity patterns within subjects [30]. Importantly, when the slow fluctuations of other frequency bands were explored (e.g., theta, alpha, beta), iEEG connectivity patterns, although observed, were anatomically coarse (extending into non-DN regions). This observation of significant, but anatomically nonspecific, iEEG correlations for slow fluctuations of lower frequency bands is consistent with earlier observations [68]. However, more recent reports, focused on different DN regions, suggest that while slow fluctuations of high-frequency activity is the most reliable correlate of resting-state fMRI connectivity, other frequency bands might also play a role, and low frequency slow fluctuations may differ between intrinsic networks [71] (Figure 3C). This work requires further exploration, and extension to other brain states where low-frequency activity is more varied.

If functional connectivity patterns truly reflect intrinsic network organization, they should exhibit some stability across brain states. Partially addressing this question, a recent study [30] investigated iEEG intrinsic connectivity between posterior nodes of the DN (PCC/RSC and AG) and found that, as anticipated, intrinsic connectivity was intact across rest, an internally directed memory task, and sleep (Figure 3B). This suggests that these slow fluctuations reflect intrinsic network properties that are subsequently modulated by behavioral state and cognition




Figure 3. Intrinsic Connectivity, Internetwork Interactions, and Electrical Stimulation of the Default Network. (A) Considerable correspondence between maps of intrinsic functional connectivity as measured with fMRI blood oxygen level-dependent (BOLD) (two leftmost panels) and intracranial electroencephalography (iEEG) power restricted to high-gamma power (HGP) (50–150 Hz) (rightmost panel) across lateral default network (DN) regions as well as other brain networks. Color indicates strength of correlated spontaneous activity at each point with a seed region in the medial prefrontal cortex (PFC) DN hub (white-rimmed circle in leftmost panel). Black circles highlight positive correlations with other default regions for both methods, and conversely, white boxes underscore anticorrelations with various non-default regions. (B) Using electrodes in the posterior cingulate cortex as seed regions, significant intrinsic connectivity is observed with the angular gyrus while performing an episodic memory task, at rest, and during sleep. (C) Spatial correlation of intrinsic connectivity maps across iEEG and fMRI BOLD in the default mode network (DMN) is driven most strongly by power in the theta (4–8 Hz) and higher frequency (50–100 Hz) ranges. Other networks [somatomotor network (SMN), dorsal attention network (DAN), and frontoparietal control network (FPC)] show similarities but also differences in their correspondence spectra. (D) Precise timing of high-frequency broadband activations in a DN hub [here, posteromedial cortex (PMC)] during an episodic memory task compared with its deactivation during math and activation of superior parietal lobule (SPL) regions during the same conditions. The delayed activation and deactivation of the PMC compared with SPL suggests that the DN lies higher in a hierarchy of information flow through the brain. (E) High-frequency electrical brain stimulation along the medial occipital and dorsomedial parietal cortical surface often elicits visual and motor effects, respectively, but has no effect within posterior parietal DN regions (posterior cingulate cortex/retrosplenial cortex). (A) Reproduced, with permission, from [70], (B) reproduced, with permission, from [30], (C) reproduced, with permission, from [71], (D) reproduced, with permission, from [104], and (E) reproduced, with permission, from [95]. Abbreviations: GSR, global signal regression; ROL, response onset latency.

[74]. Nonetheless, although these regions were correlated across all conditions, they showed the highest correlation during the self-referential episodic memory condition, during which these regions are most actively engaged (Figure 2F).

Collectively, iEEG has provided several important insights about the electrophysiology of DN intrinsic connectivity. First, a variety of methods have confirmed that the iEEG signal shows correlated slow intrinsic fluctuations (<1 Hz) between DN areas commensurate with slow spontaneous fluctuations in the fMRI BOLD signal [30,48,73,75]. Second, these patterns of iEEG intrinsic connectivity correlate well with connectivity patterns in the fMRI BOLD signal at the same sites in the same individuals [30,70,71] (Figure 3A). Third, power in specific frequency bands contributes most to slow iEEG fluctuations in intrinsic connectivity, with HFB the most consistent and anatomically specific correlate [30,70,71], although other frequency bands (<20 Hz) might also play a role [71,76] (Figure 3C). These findings are important because different frequency bands reflect distinctive neurophysiological processes: whereas changes in HFB amplitude are likely to reflect population spiking, oscillatory activity in lower frequencies might reflect neurophysiological events, such as excitability fluctuations, that coordinate long-distance communication [77]. Finally, intrinsic connectivity between DN areas is preserved across multiple cognitive states (Figure 3B), a major corroboration of findings from neuroimaging [13–16]. Going beyond mere replication, however, iEEG has shown that cross-state patterns of connectivity similarity are largely limited to high-beta and HFB power [30].

Temporal Priority and Internetwork Coordination

The millisecond temporal resolution of iEEG allows for detailed examination of the timing of DN region recruitment, as well as DN communication with other networks, providing important clues about the mechanisms of intra- and internetwork communication. For instance, there are differences in mean deactivation latencies across DN areas (Figure 2B) on the order of tens to hundreds of milliseconds [26–28], far too fast to detect with present fMRI methods. With respect to activations, although the AG and PCC/RSC show no significant differences in their mean response times [30], this might not be true for more widely dispersed regions within a network, a question that needs to be examined in future work. iEEG can also answer questions about the timing of internetwork interactions and information flow and their relation to DN activations. For instance, visual and attention networks are recruited faster than PCC and RSC during various forms of cognition (Figure 3D). Taken together, subtle timing differences suggest that DN regions might not simply activate and deactivate *en masse*, despite appearances from functional neuroimaging; and internetwork timing differences hint at hierarchies of information flow, and possibly also control, within and between intrinsic networks, each of which may have a specific ‘temporal receptive window’ related to its optimal functioning [78].

Additionally, significant phase synchronization (specifically in the 3–5 Hz theta range) occurs between RSC and more anterior regions of the MTL (outside the DN) for episodic memory judgments, but not control conditions [51]. Critically, the theta band synchronization between RSC and MTL consistently peaks prior to the onset of the relatively late (~400 ms) HFB responses in RSC, suggesting that internetwork interactions precede task-specific DN activations. These results are consistent with the finding (discussed above) that populations in the more ventral regions of the PMC, which have stronger anatomical connections to MTL regions [79,80], display earlier activations in response to episodic memory recall than populations in more dorsal aspects of the PMC [28]. Together, the iEEG evidence strongly suggests that the DN activation consistently observed during memory recall is in fact preceded by activity in, and communication with, areas outside the DN, such as anterior aspects of the MTL [51,81,82]. Collectively, these fine-scale differences in temporal priority (Figure 3D) have implications not

Box 4. Electroceutical Interventions for Clinical Dysfunctions of the Default Network

Numerous psychiatric, neurological, and neurodegenerative diseases implicate the default network (DN) [121–124]. Developing a deeper understanding of the DN's functioning in healthy people therefore lays the groundwork for better models of, and interventions for, neuropsychiatric disease. Noninvasive electroceutical interventions, like transcranial magnetic stimulation, are leading the way [125], but currently have limited ability to reach and specifically target deep brain structures (including many DN regions) in large brains, such as those of humans. As the limitations of lesioning, pharmaceuticals, and noninvasive electrical interventions for mental health disorders are increasingly appreciated [126–128], the use of intracranial electroencephalography methods for focal neuromodulation is also becoming increasingly common and sophisticated [114,129–133], and might also result in fewer long-term adverse effects [127].

Stimulation of DN regions *per se* is neither necessary for modulating DN function, nor necessarily the best means of doing so, because focal electrical stimulation has widespread effects on even distal brain regions [134,135]. Pilot studies have already investigated how focal stimulation of particular regions influences DN activity and connectivity and how these changes might relate to symptom improvements in neuropsychiatric disease such as depression and obsessive-compulsive disorder [114,115,136–138], but these studies should be viewed as valuable proofs-of-concept rather than proven therapies. Nonetheless, deep brain stimulation (DBS) interventions are reversible, adaptable, and can be continually ameliorated as both technological expertise and understanding of brain disease improve [132,133,139]. Challenges on the immediate horizon include minimizing adverse side-effects of surgical implantation and subsequent stimulation [140,141]; developing an understanding of neural mechanisms of action [134,142]; exploring whether genetics predisposes particular individuals to being more or less responsive to intracranial stimulation, as is suggested for transcranial stimulation methods [143,144]; and building 'smart' stimulation devices that act preemptively and/or adaptively [130,145].

only for our understanding of which regions initiate internally directed forms of cognition and how [83], but also for models seeking to elucidate which regions preferentially 'drive' network activity and how these can be perturbed in the service of clinical interventions [62] (Box 4).

Directionality and Time Course of Intra- and Internetwork Communication

Interactions between nodes within the DN, and between the default and other networks, can be directly investigated with iEEG using single pulse electrical stimulation (also known as CCEPs; [84]). In this approach, repeated single-pulses of electrical charge are delivered at a target electrode site while recording the responses elicited at other electrodes throughout the brain [84–87]. Research has shown that the pattern and magnitude of evoked responses predicts subject-specific intrinsic network connectivity recorded independently with resting-state fMRI [85]. Many questions about DN function and connectivity could be explored using similar paradigms to investigate both the magnitude and timing of CCEPs. For instance, several detailed explorations of the magnitude of CCEPs suggest that many bidirectional connections are asymmetrical (i.e., show differential strength) [86,88,89]. A straightforward interpretation of this asymmetry is that for any given pair of brain regions, certain regions ('projectors') are better at driving activity in the reciprocally connected brain area, whereas others ('integrators') play a more integrative or computational role [86,88,89].

A recent study went beyond investigation of the magnitude of CCEPs to examine their detailed directionality and temporal dynamics [87]. The researchers functionally identified iEEG electrode locations within subject-specific networks of interest (default, frontoparietal, and salience networks) using presurgical resting-state fMRI intrinsic connectivity in the same individuals. The CCEPs that resulted from stimulating sites within each network revealed two important findings. First, interactions among the three intrinsic networks were not mutually bidirectional: evoked responses elicited across networks were clearly asymmetrical. Second, temporal patterns of signal propagation within and between networks were not equivalent: stimulation of the frontoparietal and salience networks evoked strong responses at an earlier processing stage in all networks, whereas stimulating the DN influenced the frontoparietal and salience networks at a significantly later stage (>100 ms difference) [87]. These findings suggest differences in 'temporal receptive window' across networks [78]: the DN may be influencing other networks after they have finished their own local processing, whereas the engagement of

salience and frontoparietal networks might make a more instantaneous impact on other networks, a hypothesis needing further exploration.

Probing the Causal Importance of the Default Network

Injecting high-frequency electrical current (typically ≥ 50 Hz, ~ 1 – 10 mA) into the brain has long been known to both elicit subjective experiences [90,91] and sometimes disrupt ongoing cognitive or motor processes [92]. **High-frequency electrical brain stimulation (hf-EBS)** has been applied widely throughout cortical and subcortical structures in clinical ‘functional mapping’ sessions [93,94], but despite providing uniquely causal information about a region’s functioning, reports of hf-EBS to DN areas, especially the major hubs along the medial surface of the brain, have been scarce.

Administering more than 800 electrical stimulations throughout medial posterior brain regions in 25 passively resting epilepsy patients, our group [95] found that stimulation to PCC and RSC never yielded subjective experiences or disturbances of any kind (Figure 3E). Given the role of these areas in memory recall [96], and their selective iEEG activity both at rest and during self-referential thinking and memory recall [28–30,52], it seems likely that hf-EBS administered during internally directed tasks could cause an interruption in these cognitive functions, albeit without eliciting effects in the subjective or perceptual domain.

hf-EBS has been applied to various other putative default regions in scattered case reports [93], but virtually none of these studies has confirmed that stimulation sites were specifically within DN boundaries, or undertaken a systematic investigation including reporting of null effects. A thorough understanding of the effects of hf-EBS to DN regions is an important project for future research, and will require targeted studies employing large cohorts, precise electrode localization, and detailed subjective reports.

Relevance of the Non-Human Default Network

Although the explicit focus of this review has been on the human DN, recent years have seen the discovery of parallel (and possibly homologous) ‘default’ networks in the macaque [17], chimpanzee [97], cat [98], rat [99], and mouse [100]. A deep understanding of the human DN will need to include an account of the ways in which it is similar to potentially homologous networks in other mammals, but also what makes it unique. Here we provide a brief sketch of what is known so far about such similarities and differences, as well as highlighting how human iEEG provides unique information that goes beyond findings from animal studies.

First, functional neuroimaging methods have delineated intrinsically connected networks in animals that roughly parallel the human DN, including nodes in medial PFC, PMC, lateral parietal cortex, and lateral temporal cortex [17,97–100]. Despite the importance of these findings, cross-state comparisons have so far been sparse in animal studies because fMRI without anesthesia is challenging in most species (for an exception, see [101]). In humans, conversely, it has been straightforward to demonstrate the persistence of DN intrinsic connectivity across multiple conscious states with both fMRI and iEEG (Figure 3B [30]).

Second, task-induced deactivation of a DN region has been reported in monkeys: having macaques engage in attention or working memory tasks reliably suppressed neuron spiking in CGp [33] (a homolog of human PCC). Moreover, and reminiscent of findings in humans discussed above, less suppression of activity in CGp predicted performance errors as well as slower reaction times, again corroborating the findings from human research that DN deactivation has functional significance.

While animal studies have therefore been able, in a limited way, to study DN intrinsic connectivity (usually under anesthesia) and task-induced deactivations (in a single DN region), we are aware of no reports of task-induced activations specifically investigating DN function in the animal literature. The growing body of iEEG research investigating the many complex forms of cognition that activate DN regions in humans (such as episodic memory recall and self-referential judgments; [28–30,51,52]) will therefore provide a unique contribution to our understanding of the DN.

Concluding Remarks and Future Perspectives

Investigation of the DN has progressed rapidly since its initial discovery and has now become a major research program. The success of this project depends critically on a clear recognition of the strengths and weaknesses of the various tools at our disposal, and on their resourceful combination in research contexts. While iEEG offers spatially and temporally precise neural information at the implanted electrode sites, electrode coverage can be sparse in any given patient, and standard iEEG electrodes simultaneously record the activity of hundreds of thousands of neurons [19]. Specialized microelectrodes can resolve the differential responses of individual neurons in the human brain [102], but such methods are still employed only rarely in iEEG research.

Combining iEEG with other methods, such as fMRI, can often mitigate these concerns by providing broader context and mutually corroborative data (Box 3). And despite its limitations, human iEEG goes beyond simply replicating what is already known, or can be known, from noninvasive lines of research in humans or from invasive recordings in non-human mammalian brains. By implementing intracranial methods in human participants who can provide self-reports and execute complex cognitive-affective tasks, iEEG is making a unique contribution by illuminating what DN activation, suppression, and connectivity are (in terms of underlying electrophysiology) and what they mean (in terms of behavioral and cognitive consequences for ‘higher’ mental processes).

The major task ahead for the field is to continue moving beyond a purely correlational understanding of the DN toward high spatiotemporal resolution models of: (i) electrophysiological foundations, (ii) intraregional functional heterogeneity, (iii) intra- and internetwork communication, (iv) neural and subjective effects of direct electrical perturbation, and (iv) relationships with cognition and behavior (see Outstanding Questions). Clinicians can then leverage the knowledge gleaned from this developing neuromechanistic account to (v) better understand, and hopefully treat, neurological and psychiatric conditions involving DN dysfunction (Box 4). Advancing our understanding of the DN will therefore have many ramifications both for fathoming the marvelous complexity of the healthy brain at work, as well as treating the many ways this functioning can go awry in brain disease.

Acknowledgments

The authors are grateful to the many patients who have volunteered to participate in the studies described here, without whom this research would be impossible, as well as numerous funding agencies for their generous support. K.C.R.F. is supported by a Postdoctoral Fellowship from the Natural Sciences and Engineering Research Council (NSERC) of Canada. B.L.F. is supported by a research grant from the U.S. National Institute of Mental Health (R00MH103479). A.K. is supported by a Banting Postdoctoral Fellowship from the Canadian Institutes of Health Research (CIHR). A.L.D. is supported by a postdoctoral fellowship from the National Institute of Child Health and Human Development (1F32HD087028-01). J.P. is supported by research grants from the U.S. National Institute of Neurological Disorders and Stroke (R01NS078396), U.S. National Institute of Mental Health (1R01MH109954-01), and the U.S. National Science Foundation (BCS1358907).

Outstanding Questions

How is ‘deactivation’ instantiated at the single-cell level in the default network (DN) and elsewhere? Can single-neuron studies in humans clarify whether reduced single-neuron activity (i.e., firing rate) plays any functional role and whether particular neuronal types (e.g., interneurons) are involved?

What is the functional significance of the rapid (<300 ms) responses in DN regions during the switch from engagement in externally directed tasks to rest? What neural mechanisms are involved? Is this ‘switch signal’ the cause or the result of disengagement from externally oriented tasks?

What is the significance of functionally heterogeneous neuronal populations within other DN regions beyond the posteromedial cortex (PMC)?

What is the electrophysiological basis of DN involvement in behavioral correlates such as attentional stability, tracking changing narratives, and functioning on ‘autopilot’?

How are both competitive and cooperative internetwork interactions mediated and instantiated at the electrophysiological level, both at rest and during tasks? Are specific frequency ranges of electrophysiological activity of particular importance?

What mechanisms drive the slow dynamics of DN intrinsic connectivity? Is this slow variation causally linked to other important psychophysiological processes? Are glial cells important?

What are the subjective effects of direct electrical stimulation to DN areas beyond the PMC? Does stimulation affect or impair performance on externally and/or internally directed tasks?

How do putatively homologous ‘default’ networks in non-human primates and rodents differ functionally and electrophysiologically from the human DN?

How can iEEG methods best be harnessed to understand and treat the

References

1. Roberts, R.M. (1989) *Serendipity: Accidental Discoveries in Science*, Wiley
2. Kuhn, T.S. (1975) *The Structure of Scientific Revolutions*. (2nd edn), University of Chicago Press
3. Buckner, R.L. (2012) The serendipitous discovery of the brain's default network. *Neuroimage* 62, 1137–1145
4. Raichle, M.E. *et al.* (2001) A default mode of brain function. *Proc. Natl. Acad. Sci. U. S. A.* 98, 678–682
5. Shulman, G.L. *et al.* (1997) Common blood flow changes across visual tasks: II. Decreases in cerebral cortex. *J. Cogn. Neurosci.* 9, 648–663
6. Raichle, M.E. (2009) A paradigm shift in functional brain imaging. *J. Neurosci.* 29, 12729–12734
7. Raichle, M.E. (2010) Two views of brain function. *Trends Cogn. Sci.* 14, 180–190
8. Gusnard, D.A. and Raichle, M.E. (2001) Searching for a baseline: functional imaging and the resting human brain. *Nat. Rev. Neurosci.* 2, 685
9. Yeo, B.T.T. *et al.* (2011) The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J. Neurophysiol.* 106, 1125–1165
10. Beckmann, C.F. *et al.* (2005) Investigations into resting-state connectivity using independent component analysis. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 360, 1001–1013
11. Damoiseaux, J. *et al.* (2006) Consistent resting-state networks across healthy subjects. *Proc. Natl. Acad. Sci.* 103, 13848–13853
12. Greicius, M.D. *et al.* (2003) Functional connectivity in the resting brain: a network analysis of the default mode hypothesis. *Proc. Natl. Acad. Sci.* 100, 253–258
13. Cole, M.W. *et al.* (2014) Intrinsic and task-evoked network architectures of the human brain. *Neuron* 83, 238–251
14. Krienen, F.M. *et al.* (2014) Reconfigurable task-dependent functional coupling modes cluster around a core functional architecture. *Philos. Trans. R. Soc. B* 369, 20130526
15. Larson-Prior, L.J. *et al.* (2009) Cortical network functional connectivity in the descent to sleep. *Proc. Natl. Acad. Sci.* 106, 4489–4494
16. Vanhaudenhuyse, A. *et al.* (2010) Default network connectivity reflects the level of consciousness in non-communicative brain-damaged patients. *Brain* 133, 161–171
17. Vincent, J.L. *et al.* (2007) Intrinsic functional architecture in the anesthetized monkey brain. *Nature* 447, 83–86
18. Raichle, M.E. (2015) The brain's default mode network. *Annu. Rev. Neurosci.* 38, 433–447
19. Parvizi, J. and Kastner, S. Promises and limitations of human intracranial electroencephalography. *Nat. Neurosci.* (in press). <https://doi.org/10.1038/s41593-018-0108-2>.
20. Logothetis, N.K. *et al.* (2001) Neurophysiological investigation of the basis of the fMRI signal. *Nature* 412, 150–157
21. Mukamel, R. *et al.* (2005) Coupling between neuronal firing, field potentials, and fMRI in human auditory cortex. *Science* 309, 951–954
22. Nir, Y. *et al.* (2007) Coupling between neuronal firing rate, gamma LFP, and BOLD fMRI is related to interneuronal correlations. *Curr. Biol.* 17, 1275–1285
23. Lachaux, J.-P. *et al.* (2008) Silence is golden: transient neural deactivation in the prefrontal cortex during attentive reading. *Cereb. Cortex* 18, 443–450
24. Jung, J. *et al.* (2010) Brain responses to success and failure: direct recordings from human cerebral cortex. *Hum. Brain Mapp.* 31, 1217–1232
25. Miller, K.J. *et al.* (2009) Direct electrophysiological measurement of human default network areas. *Proc. Natl. Acad. Sci.* 106, 12174–12177
26. Jerbi, K. *et al.* (2010) Exploring the electrophysiological correlates of the default-mode network with intracerebral EEG. *Front. Syst. Neurosci.* 4, 27
27. Ossandón, T. *et al.* (2011) Transient suppression of broadband gamma power in the default-mode network is correlated with task complexity and subject performance. *J. Neurosci.* 31, 14521–14530
28. Foster, B.L. *et al.* (2012) Neural populations in human posteromedial cortex display opposing responses during memory and numerical processing. *Proc. Natl. Acad. Sci.* 109, 15514–15519
29. Dastjerdi, M. *et al.* (2011) Differential electrophysiological response during rest, self-referential, and non-self-referential tasks in human posteromedial cortex. *Proc. Natl. Acad. Sci.* 108, 3023–3028
30. Foster, B.L. *et al.* (2015) Intrinsic and task-dependent coupling of neuronal population activity in human parietal cortex. *Neuron* 86, 578–590
31. Birn, R.M. *et al.* (2008) The effect of respiration variations on independent component analysis results of resting state functional connectivity. *Hum. Brain Mapp.* 29, 740–750
32. Shmuel, A. *et al.* (2002) Sustained negative BOLD, blood flow and oxygen consumption response and its coupling to the positive response in the human brain. *Neuron* 36, 1195–1210
33. Hayden, B.Y. *et al.* (2009) Electrophysiological correlates of default-mode processing in macaque posterior cingulate cortex. *Proc. Natl. Acad. Sci.* 106, 5948–5953
34. Ramot, M. *et al.* (2012) A widely distributed spectral signature of task-negative electrocorticography responses revealed during a visuomotor task in the human cortex. *J. Neurosci.* 32, 10458–10469
35. Daitch, A.L. and Parvizi, J. (2017) Heterogeneous memory- and rest-related activity within the human posteromedial cortex. *Annual Meeting of the Organization for Human Brain Mapping*
36. Weissman, D. *et al.* (2006) The neural bases of momentary lapses in attention. *Nat. Neurosci.* 9, 971–978
37. McKiernan, K.A. *et al.* (2003) A parametric manipulation of factors affecting task-induced deactivation in functional neuroimaging. *J. Cogn. Neurosci.* 15, 394–408
38. Mason, M.F. *et al.* (2007) Wandering minds: the default network and stimulus-independent thought. *Science* 315, 393–395
39. Teasdale, J.D. *et al.* (1995) Stimulus-independent thought depends on central executive resources. *Memory Cogn.* 23, 551–559
40. Dixon, M.L. *et al.* (2014) A framework for understanding the relationship between externally and internally directed cognition. *Neuropsychologia* 62, 321–330
41. Buckner, R.L. *et al.* (2008) The brain's default network: anatomy, function, and relevance to disease. *Ann. N. Y. Acad. Sci.* 1124, 1–38
42. Gusnard, D.A. *et al.* (2001) Medial prefrontal cortex and self-referential mental activity: relation to a default mode of brain function. *Proc. Natl. Acad. Sci. U. S. A.* 98, 4259–4264
43. Fox, K.C.R. *et al.* (2015) The wandering brain: meta-analysis of functional neuroimaging studies of mind-wandering and related spontaneous thought processes. *Neuroimage* 111, 611–621
44. Christoff, K. *et al.* (2009) Experience sampling during fMRI reveals default network and executive system contributions to mind wandering. *Proc. Natl. Acad. Sci. U. S. A.* 106, 8719–8724
45. Addis, D.R. *et al.* (2007) Remembering the past and imagining the future: common and distinct neural substrates during event construction and elaboration. *Neuropsychologia* 45, 1363–1377
46. Svoboda, E. *et al.* (2006) The functional neuroanatomy of autobiographical memory: a meta-analysis. *Neuropsychologia* 44, 2189–2208
47. Miller, K.J. *et al.* (2007) Spectral changes in cortical surface potentials during motor movement. *J. Neurosci.* 27, 2424–2432
48. Foster, B.L. and Parvizi, J. (2012) Resting oscillations and cross-frequency coupling in the human posteromedial cortex. *Neuroimage* 60, 384–391

many neuropsychiatric disorders that implicate the DN? What effect does focal neuromodulation have on the wider dynamics of the default and other brain networks?

What are the electrophysiological dynamics and functional role of sub-cortical components of the DN, such as the thalamus and cerebellum?

49. Spreng, R.N. *et al.* (2009) The common neural basis of autobiographical memory, prospection, navigation, theory of mind, and the default mode: a quantitative meta-analysis. *J. Cogn. Neurosci.* 21, 489–510
50. Andrews-Hanna, J.R. *et al.* (2014) The default network and self-generated thought: component processes and dynamic control. *Ann. N. Y. Acad. Sci.* 1316, 29–52
51. Foster, B.L. *et al.* (2013) Human retrosplenial cortex displays transient theta phase locking with medial temporal cortex prior to activation during autobiographical memory retrieval. *J. Neurosci.* 33, 10439–10446
52. Daitch, A.L. and Parvizi, J. Activations and deactivations in different subregions of the human posteromedial cortex: heterogeneity in space and time. *Proc. Natl. Acad. Sci.* (in press)
53. Northoff, G. *et al.* (2006) Self-referential processing in our brain—a meta-analysis of imaging studies on the self. *Neuroimage* 31, 440–457
54. Andrews-Hanna, J.R. *et al.* (2010) Functional-anatomic fractionation of the brain's default network. *Neuron* 65, 550–562
55. Braga, R.M. and Buckner, R.L. (2017) Parallel interdigitated distributed networks within the individual estimated by intrinsic functional connectivity. *Neuron* 95, 457–471
56. Leech, R. *et al.* (2011) Fractionating the default mode network: distinct contributions of the ventral and dorsal posterior cingulate cortex to cognitive control. *J. Neurosci.* 31, 3217–3224
57. Margulies, D.S. *et al.* (2007) Mapping the functional connectivity of anterior cingulate cortex. *Neuroimage* 37, 579–588
58. Crittenden, B.M. *et al.* (2015) Recruitment of the default mode network during a demanding act of executive control. *Elife* 4, e06481
59. Hayden, B.Y. *et al.* (2010) Cognitive control signals in posterior cingulate cortex. *Front. Hum. Neurosci.* 4, 223
60. Ellamil, M. *et al.* (2016) Dynamics of neural recruitment surrounding the spontaneous arising of thoughts in experienced mindfulness practitioners. *Neuroimage* 136, 186–196
61. Sporns, O. (2014) Contributions and challenges for network models in cognitive neuroscience. *Nat. Neurosci.* 17, 652–660
62. Bassett, D.S. and Sporns, O. (2017) Network neuroscience. *Nat. Neurosci.* 20, 353
63. Adrian, E.D. and Matthews, B.H. (1934) The Berger rhythm: potential changes from the occipital lobes in man. *Brain* 57, 355–385
64. Berger, H. (1929) Über das elektroencephalogramm des menschen. *Eur. Arch. Psychiatry Clin. Neurosci.* 87, 527–570 (in German)
65. Buzsáki, G. (2002) Theta oscillations in the hippocampus. *Neuron* 33, 325–340
66. Biswal, B. *et al.* (1995) Functional connectivity in the motor cortex of resting human brain using echo-planar MRI. *Magn. Reson. Med.* 34, 537–541
67. Buckner, R.L. *et al.* (2013) Opportunities and limitations of intrinsic functional connectivity MRI. *Nat. Neurosci.* 16, 832–837
68. Nir, Y. *et al.* (2008) Interhemispheric correlations of slow spontaneous neuronal fluctuations revealed in human sensory cortex. *Nat. Neurosci.* 11, 1100–1108
69. Leopold, D.A. *et al.* (2003) Very slow activity fluctuations in monkey visual cortex: implications for functional brain imaging. *Cereb. Cortex* 13, 422–433
70. Keller, C.J. *et al.* (2013) Neurophysiological investigation of spontaneous correlated and anticorrelated fluctuations of the BOLD signal. *J. Neurosci.* 33, 6333–6342
71. Hacker, C.D. *et al.* (2017) Frequency-specific electrophysiologic correlates of resting state fMRI networks. *Neuroimage* 149, 446–457
72. Kucyi, A. *et al.* (2017) Consistent ECoG-fMRI Correspondence of Intrinsic Networks Across fMRI Denoising Strategies, 47th Annual Meeting of the Society for Neuroscience, Washington, D.C.
73. Ko, A.L. *et al.* (2011) Quasi-periodic fluctuations in default mode network electrophysiology. *J. Neurosci.* 31, 11728–11732
74. Buzsáki, G. (2006) *Rhythms of the Brain*, Oxford University Press
75. Ko, A.L. *et al.* (2013) Identifying functional networks using endogenous connectivity in gamma band electrocorticography. *Brain Connect.* 3, 491–502
76. Wang, L. *et al.* (2012) Electrophysiological low-frequency coherence and cross-frequency coupling contribute to BOLD connectivity. *Neuron* 76, 1010–1020
77. Fries, P. (2005) A mechanism for cognitive dynamics: neuronal communication through neuronal coherence. *Trends Cogn. Sci.* 9, 474–480
78. Honey, C.J. *et al.* (2012) Slow cortical dynamics and the accumulation of information over long timescales. *Neuron* 76, 423–434
79. Kobayashi, Y. and Amaral, D.G. (2003) Macaque monkey retrosplenial cortex: II. Cortical afferents. *J. Comp. Neurol.* 466, 48–79
80. Parvizi, J. *et al.* (2006) Neural connections of the posteromedial cortex in the macaque. *Proc. Natl. Acad. Sci.* 103, 1563–1568
81. Burke, J.F. *et al.* (2014) Theta and high-frequency activity mark spontaneous recall of episodic memories. *J. Neurosci.* 34, 11355–11365
82. Gelbard-Sagiv, H. *et al.* (2008) Internally generated reactivation of single neurons in human hippocampus during free recall. *Science* 322, 96–101
83. Fox, K.C.R. *et al.* (2016) The neurobiology of self-generated thought from cells to systems: integrating evidence from lesion studies, human intracranial electrophysiology, neurochemistry, and neuroendocrinology. *Neuroscience* 335, 134–150
84. Matsumoto, R. *et al.* (2004) Functional connectivity in the human language system: a cortico-cortical evoked potential study. *Brain* 127, 2316–2330
85. Keller, C.J. *et al.* (2011) Intrinsic functional architecture predicts electrically evoked responses in the human brain. *Proc. Natl. Acad. Sci.* 108, 10308–10313
86. Keller, C.J. *et al.* (2014) Corticocortical evoked potentials reveal projectors and integrators in human brain networks. *J. Neurosci.* 34, 9152–9163
87. Shine, J.M. *et al.* (2017) Distinct patterns of temporal and directional connectivity among intrinsic networks in the human brain. *J. Neurosci.* 37, 9667–9674
88. Entz, L. *et al.* (2014) Evoked effective connectivity of the human neocortex. *Hum. Brain Mapp.* 35, 5736–5753
89. Keller, C.J. *et al.* (2014) Mapping human brain networks with cortico-cortical evoked potentials. *Philos. Trans. R. Soc. B* 369, 20130528
90. Bartholow, R. (1874) Art. I. Experimental investigations into the functions of the human brain. *Am. J. Med. Sci.* 66, 305–313
91. Penfield, W. and Boldrey, E. (1937) Somatic motor and sensory representation in the cerebral cortex of man as studied by electrical stimulation. *Brain* 60, 389–443
92. Ojemann, G. *et al.* (1989) Cortical language localization in left, dominant hemisphere: an electrical stimulation mapping investigation in 117 patients. *J. Neurosurg.* 71, 316–326
93. Selimbeyoglu, A. and Parvizi, J. (2010) Electrical stimulation of the human brain: perceptual and behavioral phenomena reported in the old and new literature. *Front. Hum. Neurosci.* 4, 46
94. Borchers, S. *et al.* (2012) Direct electrical stimulation of human cortex—the gold standard for mapping brain functions? *Nat. Rev. Neurosci.* 13, 63
95. Foster, B.L. and Parvizi, J. (2017) Direct cortical stimulation of human posteromedial cortex. *Neurology* 88, 685–691
96. Shannon, B.J. and Buckner, R.L. (2004) Functional-anatomic correlates of memory retrieval that suggest nontraditional processing roles for multiple distinct regions within posterior parietal cortex. *J. Neurosci.* 24, 10084–10092
97. Rilling, J.K. *et al.* (2007) A comparison of resting-state brain activity in humans and chimpanzees. *Proc. Natl. Acad. Sci.* 104, 17146–17151

98. Popa, D. *et al.* (2009) Contrasting activity profile of two distributed cortical networks as a function of attentional demands. *J. Neurosci.* 29, 1191–1201
99. Lu, H. *et al.* (2012) Rat brains also have a default mode network. *Proc. Natl. Acad. Sci.* 109, 3979–3984
100. Stafford, J.M. *et al.* (2014) Large-scale topology and the default mode network in the mouse connectome. *Proc. Natl. Acad. Sci.* 111, 18745–18750
101. Barttfeld, P. *et al.* (2015) Signature of consciousness in the dynamics of resting-state brain activity. *Proc. Natl. Acad. Sci.* 112, 887–892
102. Rutishauser, U. *et al.* (2018) Single-neuron representation of memory strength and recognition confidence in left human posterior parietal cortex. *Neuron* 97, 209–220
103. Jacobs, J. and Kahana, M.J. (2010) Direct brain recordings fuel advances in cognitive electrophysiology. *Trends Cogn. Sci.* 14, 162–171
104. Raccach, O. *et al.* (2017) *Relative Timing of Opposing Responses in Default Mode and Dorsal Attention Networks*, Annual Meeting of the Organization for Human Brain Mapping, Vancouver, B.C., Canada
105. Greicius, M.D. *et al.* (2008) Resting-state functional connectivity reflects structural connectivity in the default mode network. *Cereb. Cortex* 19, 72–78
106. Buzsáki, G. *et al.* (2012) The origin of extracellular fields and currents—EEG, ECoG, LFP and spikes. *Nat. Rev. Neurosci.* 13, 407–420
107. Yuste, R. (2015) From the neuron doctrine to neural networks. *Nat. Rev. Neurosci.* 16, 487–497
108. Mueller, S. *et al.* (2013) Individual variability in functional connectivity architecture of the human brain. *Neuron* 77, 586–595
109. Gordon, E.M. *et al.* (2017) Individual variability of the system-level organization of the human brain. *Cereb. Cortex* 27, 386–399
110. Laumann, T.O. *et al.* (2015) Functional system and areal organization of a highly sampled individual human brain. *Neuron* 87, 657–670
111. Wang, D. *et al.* (2015) Parcellating cortical functional networks in individuals. *Nat. Neurosci.* 18, 1853
112. Parvizi, J. *et al.* (2013) The will to persevere induced by electrical stimulation of the human cingulate gyrus. *Neuron* 80, 1359–1367
113. Figeo, M. *et al.* (2013) Deep brain stimulation restores frontostriatal network activity in obsessive-compulsive disorder. *Nat. Neurosci.* 16, 386–387
114. Mayberg, H.S. *et al.* (2005) Deep brain stimulation for treatment-resistant depression. *Neuron* 45, 651–660
115. Lozano, A.M. *et al.* (2008) Subcallosal cingulate gyrus deep brain stimulation for treatment-resistant depression. *Biol. Psychiatry* 64, 461–467
116. Carmichael, D.W. *et al.* (2007) Functional MRI. with active, fully implanted, deep brain stimulation systems: safety and experimental confounds. *Neuroimage* 37, 508–517
117. Carmichael, D.W. *et al.* (2010) Feasibility of simultaneous intracranial EEG-fMRI in humans: a safety study. *Neuroimage* 49, 379–390
118. Dalal, S.S. *et al.* (2009) Simultaneous MEG and intracranial EEG recordings during attentive reading. *Neuroimage* 45, 1289–1304
119. Dubarry, A.-S. *et al.* (2014) Simultaneous recording of MEG, EEG and intracerebral EEG during visual stimulation: from feasibility to single-trial analysis. *Neuroimage* 99, 548–558
120. Logothetis, N.K. *et al.* (2010) The effects of electrical microstimulation on cortical signal propagation. *Nat. Neurosci.* 13, 1283–1291
121. Christoff, K. *et al.* (2016) Mind-wandering as spontaneous thought: a dynamic framework. *Nat. Rev. Neurosci.* 17, 718–731
122. Whitfield-Gabrieli, S. and Ford, J.M. (2012) Default mode network activity and connectivity in psychopathology. *Annu. Rev. Clin. Psychol.* 8, 49–76
123. Kaiser, R.H. *et al.* (2015) Large-scale network dysfunction in major depressive disorder: a meta-analysis of resting-state functional connectivity. *JAMA Psychiatry* 72, 603–611
124. Anticevic, A. *et al.* (2012) The role of default network deactivation in cognition and disease. *Trends Cogn. Sci.* 16, 584–592
125. Kuo, M.-F. *et al.* (2014) Therapeutic effects of non-invasive brain stimulation with direct currents (tDCS) in neuropsychiatric diseases. *Neuroimage* 85, 948–960
126. Underwood, E. (2016) Cadaver study casts doubts on how zapping brain may boost mood, relieve pain. Published online April 20, 2016. <https://doi.org/10.1126/science.aaf9938>
127. Kern, D.S. and Kumar, R. (2007) Deep brain stimulation. *Neurologist* 13, 237–252
128. Cipriani, A. *et al.* (2009) Comparative efficacy and acceptability of 12 new-generation antidepressants: a multiple-treatments meta-analysis. *Lancet* 373, 746–758
129. Deuschl, G. *et al.* (2006) A randomized trial of deep-brain stimulation for Parkinson's disease. *N. Engl. J. Med.* 355, 896–908
130. Fischell, R.E. *et al.* (2000) NeuroPace Inc. System for treatment of neurological disorders, US 6016449
131. Holtzheimer, P.E. and Mayberg, H.S. (2011) Deep brain stimulation for psychiatric disorders. *Annu. Rev. Neurosci.* 34, 289–307
132. De Ridder, D. *et al.* (2017) State of the art: novel applications for cortical stimulation. *Neuromodulation* 20, 206–214
133. Roy, H.A. *et al.* (2018) State of the art: novel applications for deep brain stimulation. *Neuromodulation* 21, 126–134
134. McIntyre, C.C. and Hahn, P.J. (2010) Network perspectives on the mechanisms of deep brain stimulation. *Neurobiol. Dis.* 38, 329–337
135. Alhourani, A. *et al.* (2015) Network effects of deep brain stimulation. *J. Neurophysiol.* 114, 2105–2117
136. Figeo, M. *et al.* (2011) Dysfunctional reward circuitry in obsessive-compulsive disorder. *Biol. Psychiatry* 69, 867–874
137. Menzies, L. *et al.* (2008) Integrating evidence from neuroimaging and neuropsychological studies of obsessive-compulsive disorder: the orbitofronto-striatal model revisited. *Neurosci. Biobehav. Rev.* 32, 525–549
138. Harrison, B.J. *et al.* (2009) Altered corticostriatal functional connectivity in obsessive-compulsive disorder. *Arch. Gen. Psychiatry* 66, 1189–1200
139. Morishita, T. *et al.* (2014) Deep brain stimulation for treatment-resistant depression: systematic review of clinical outcomes. *Neurotherapeutics* 11, 475–484
140. Kisely, S. *et al.* (2014) Deep brain stimulation for obsessive-compulsive disorder: a systematic review and meta-analysis. *Psychol. Med.* 44, 3533–3542
141. Saleh, C. and Fontaine, D. (2015) Deep brain stimulation for psychiatric diseases: what are the risks? *Curr. Psychiatry Rep.* 17, 33
142. Fox, M.D. *et al.* (2014) Resting-state networks link invasive and noninvasive brain stimulation across diverse psychiatric and neurological diseases. *Proc. Natl. Acad. Sci.* 111, E4367–E4375
143. Chhabra, H. *et al.* (2016) Transcranial direct current stimulation and neuroplasticity genes: implications for psychiatric disorders. *Acta Neuropsychiatrica* 28, 1–10
144. Cheeran, B. *et al.* (2008) A common polymorphism in the brain-derived neurotrophic factor gene (BDNF) modulates human cortical plasticity and the response to rTMS. *J. Physiol.* 586, 5717–5725
145. De Ridder, D. and Vanneste, S. (2016) Visions on the future of medical devices in spinal cord stimulation: what medical device is needed? *Expert Rev. Med. Devices* 13, 233–242

146. Miller, K.J. (2010) Broadband spectral change: evidence for a macroscale correlate of population firing rate? *J. Neurosci.* 30, 6477–6479
147. Winawer, J. *et al.* (2013) Asynchronous broadband signals are the principal source of the BOLD response in human visual cortex. *Curr. Biol.* 23, 1145–1153
148. Jiruska, P. *et al.* (2017) Update on the mechanisms and roles of high-frequency oscillations in seizures and epileptic disorders. *Epilepsia* 58, 1330–1339
149. Schevon, C.A. *et al.* (2009) Spatial characterization of interictal high frequency oscillations in epileptic neocortex. *Brain* 132, 3047–3059
150. Ray, S. *et al.* (2008) Neural correlates of high-gamma oscillations (60–200 Hz) in macaque local field potentials and their potential implications in electrocorticography. *J. Neurosci.* 28, 11526–11536
151. Ray, S. and Maunsell, J.H. (2011) Different origins of gamma rhythm and high-gamma activity in macaque visual cortex. *PLoS Biol.* 9, e1000610
152. Liu, J. and Newsome, W.T. (2006) Local field potential in cortical area MT: stimulus tuning and behavioral correlations. *J. Neurosci.* 26, 7779–7790
153. Kreiman, G. *et al.* (2006) Object selectivity of local field potentials and spikes in the macaque inferior temporal cortex. *Neuron* 49, 433–445
154. Manning, J.R. *et al.* (2009) Broadband shifts in local field potential power spectra are correlated with single-neuron spiking in humans. *J. Neurosci.* 29, 13613–13620
155. Niessing, J. *et al.* (2005) Hemodynamic signals correlate tightly with synchronized gamma oscillations. *Science* 309, 948–951
156. Flinker, A. *et al.* (2011) Sub-centimeter language organization in the human temporal lobe. *Brain Lang.* 117, 103–109
157. Mesgarani, N. *et al.* (2014) Phonetic feature encoding in human superior temporal gyrus. *Science* 348, 1006–1010
158. Hermes, D. *et al.* (2012) Neurophysiologic correlates of fMRI in human motor cortex. *Hum. Brain Mapp.* 33, 1689–1699
159. Canolty, R.T. *et al.* (2006) High gamma power is phase-locked to theta oscillations in human neocortex. *Science* 313, 1626–1628
160. Park, S.H. *et al.* (2017) Functional subpopulations of neurons in a macaque face patch revealed by single-unit fMRI mapping. *Neuron* 95, 971–981
161. Katzner, S. *et al.* (2009) Local origin of field potentials in visual cortex. *Neuron* 61, 35–41
162. Engel, A.K. *et al.* (1990) Stimulus-dependent neuronal oscillations in cat visual cortex: inter-columnar interaction as determined by cross-correlation analysis. *Eur. J. Neurosci.* 2, 588–606
163. Kucyi, A. *et al.* (2016) Spontaneous default network activity reflects behavioral variability independent of mind-wandering. *Proc. Natl. Acad. Sci.* 113, 13899–13904